

Sample Analyzed for DOMVR in the Commission II of the ICCP



Insert a microscopy image of the sample with graphic scale



Insert the distribution of vitrinite reflectance means

nthe	Commiss		the				Co,		RGA
	Sample Information						IANO.		NIC P
	Code: OM0	C3A	Year	of A	Analysis:	2009	TIN NO.		, ,
	Type of Sam	ple: Outcrop							
	Location and	Location and/or Fm.: Asturias Mesozoic Cover. Rodiles Fm							
	Country:	Spain			Age (Perio	d): Pier	nsbachia	n	-
	Dep. Enviror	nment: Marine							-
	Coordinates	Long.		Co	ordinates L	.at.			-
ICCP WG:	Organic Matter Cond	centration							
Convenor:	João Graciano Mendonça Filho E-mail: graciano@					geologia.ufrj.br			
Exercise I	nformation				(Other o	data Ava	ailable	
Report: Mendonça Filho 2009 Participants N: 14						🗙 Chei 🗙 Rock	mical Ana Eval	alysis	
Group Mean (VRr%) 1.04 Group Stdv.: 0.112						Spectral Fluorescence			
Averaged Unsigned Multiple Stdv.: 0.891							chemistry ges availa		
Coef. of Variation: 10.7 Scattering Index: 0			0.8	B Others (indicate comments)				ate in	

FOR CO

Comments:

Exercise focused on the effect of concentration procedures on the vitrinite reflectance and other optical parameters. See also Mendonça Filho et al. (International Journal of Coal Geology 84 (2010) 154–165). Difficulties to distinguish primary vitrinite and bitumen

Sample Analyzed for DOMVR in the Commission II of the ICCP LEGEND

Code: refers to the sample code as distributed for the round robin exercises

ICCP WG: name of the WG in which the exercise was run

Microscopy images: Please indicate in the image as much information as possible regarding illumination conditions and identification of components. If you use fluorescence and white light images. Insert them as a single image.

Histogram with reflectance readings: Please build up an histogram with the individual vitrinite reflectance means reported by participants to represent the scatter of the readings in the exercise

Report: indicate the name of the report in which the results of this sample are available as recorded in the webpage (i.e. Bostick 1982; Borrego 2006, etc...)

Participants N: number of results included in the exercise

Group mean (VRr %): refers to the group mean resulting of averaging the individual mean reflectance values reported by participants.

Group Stdv: refers to the group standard deviation resulting from the individual mean reflectance values reported by participants.

<u>Averaged Unsigned Multiple Stdv.</u>: refers to the Average value of the individual Unsigned Multiple of the Standard Deviations, calculated for each participant against the group mean and group standard deviation data. This statistical is used in the ICCP Accreditation Programms to assess the precision of the participants. Average Unsigned Multiple Stdv.=Summa(absolute value [participant VRr-Group Mean)/Group Stdv.])

<u>Coefficient of Variation</u>: allows comparing the dispersion of results regardless the value of the mean. Coefficient of variation=Group Stdv *100/ Group Mean.

<u>Scattering Index</u>: allows an estimation of the reliability of the values based on the Coefficient of Variation and the number of participants. Scattering Index=Coefficient of Variation/N of participants

<u>Comments</u>: Please indicate whatever information you consider relevant. Information to include is: objectives of the working group, indication about fluorescence properties, abundance of vitrinite particles to be measured, difficulties in sample preparation or polishing, possibility of suppressed values, the main conclusions about the characterization of the samples.